

OXYFLY®

Product Label

Disclaimer

This information summarizes the current status of our knowledge and may differ from the legally binding, country specific use instructions provided with the product. Novartis Animal Health Inc. is not liable for damages or injury caused by the use of its products different from those specified on the pack label.

Composition

100 g/l (9,62 % w/w) lambda-cyhalothrin as a microencapsulated suspension in water.

Direction for use

Application rates

Target	Amount of product per 5 liters water (ml)	Area treated (m ²)	Concentration (mg a.i./m ²)
Flies	50	200	25
Other insects	50	As crack and crevice treatment	25

Oxyfly Precautions / Statutory Conditions Relating to Use

- WASH ANY CONTAMINATION from skin or eyes immediately
- WASH HANDS AND EXPOSED SKIN before meals and after use
- Keep only in the original container
- Keep container tightly closed
- KEEP IN A SAFE PLACE
- Keep out of reach of children
- Keep away from food drink and animal feeding stuffs
- COLLECT EGGS, AND REMOVE FEED, EXPOSED WATER AND MILK BEFORE APPLICATION

Conditions

Store in original, tightly closed container away from sun and damp in a well ventilated area under lock and key. Avoid storage below 35oC. Protect from frost.

AVOID EXCESSIVE CONTAMINATION OF COVERALLS AND LAUNDRER
REGULARLY

Do not breathe spray mist.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

DO NOT CONTAMINATE FOODSTUFFS, EATING UTENSILS OR FOOD CONTACT SURFACES other than storage boxes or containers.

COVER ALL WATER STORAGE TANKS before application.

This material and its container must be disposed of in a safe way.

When using do not eat drink or smoke.

Avoid contact with skin.

FOR INDOOR USE ONLY.

UNPROTECTED PERSONS AND ANIMALS SHOULD BE KEPT AWAY FROM TREATED AREAS UNTIL SURFACES ARE DRY.

DO NOT APPLY DIRECTLY TO ANIMALS.

REMOVE OR COVER ALL FISH TANKS AND BOWLS before application.

DO NOT APPLY DIRECTLY TO LIVESTOCK AND POULTRY.

EXCLUDE ALL PERSONS AND ANIMALS DURING TREATMENT.

Protect milk machinery and containers from contamination.

APPLY ONLY IN AREAS INACCESSIBLE TO LIVESTOCK AND POULTRY.

READ ALL PRECAUTIONS BEFORE USE.